

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 July 2002 (25.07.2002)

PCT

(10) International Publication Number
WO 02/057797 A2

(51) International Patent Classification⁷: **G01N 33/82**

(21) International Application Number: **PCT/US01/47267**

(22) International Filing Date: 3 December 2001 (03.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: *PENDING*
09/761,969 16 January 2001 (16.01.2001) US

(71) Applicant: **QUEST DIAGNOSTICS INC** [US/US];
33608 Ortega Highway, San Juan Capistrano, CA 92690
(US).

(72) Inventors: **GARRITY, Martha**; 151 Avenida Princessa,
San Clemente, CA 92672 (US). **TRAN, Jacqueline**; 8401
Conner Circle, Westminster, CA 92683 (US).

(74) Agent: **STOUT, Donald, E.**; Stout, Uxa, Buyan &
Mullins, 4 Venture, Suite 300, Irvine, CA 92618 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 02/057797 A2

(54) Title: **VITAMIN D ASSAY**

(57) Abstract: The present invention features a kit and a method of using the kit for determining a concentration of a vitamin D component. The kit comprises a releasing composition and a detecting composition. The releasing composition comprises an aqueous base component. In one embodiment, the releasing composition is substantially free from an organic solvent.

VITAMIN D ASSAY

5 Background of the Invention

In 1928 Adolf Windaus was awarded the Nobel Prize in chemistry for his research on substances of significant biological importance. One of these substances was Vitamin D. Since then, much more is known about vitamin
10 D, its chemistry and its pharmacological effects and dynamics.

Vitamin D is not widely found in food, but rather, it is produced by the skin. Once produced, it undergoes 25-hydroxylation to form a 25-hydroxy-D (25-OH-D) in the
15 liver. The circulating concentration of 25-OH-D is considered to be an important indicator of vitamin D status in man. For example, hypovitaminosis, which results from the insufficient endogenous production of vitamin D in the skin, and insufficient dietary
20 supplementation, and/or inability of the small intestine to absorb adequate amounts of vitamin D from dietary sources, results in hypocalcemia and hypophosphatemia and corresponding secondary hyperparathyroidism. Hypovitaminosis D also results in disturbances in mineral
25 metabolism (i.e., rickets and osteomalacia in children and adults, respectively).

Serum 25-OH-D-levels are also found to be lower than normal in intestinal malabsorption syndromes, liver disorders (chronic and acute), and nephrotic syndromes.
30 In osteopenia, especially in the aged, serum 25-OH-D levels are often found to be lower than normal. In cases of vitamin D intoxication, serum 25-OH-D level is found,

as expected, to be higher than normal.

Once hydroxylated, 25-OH-D is again hydroxylated in the kidney to give the hormonal form 1,25-dihydroxy-vitamin D ($1,25-(OH)_2-D$). The $1,25-(OH)_2-D$ level in
5 blood is also an important indicator of certain diseases. For example, a low level of $1,25-(OH)_2-D$ is indicative of kidney failure and/or osteoporosis.

Considering their pathological importance, tremendous efforts have been directed towards developing
10 assays for accurately measuring concentrations of 25-OH-D and/or $1,25-(OH)_2-D$ in circulation.

For example, 25-OH-D concentrations in blood have been measured by high performance liquid chromatography (HPLC) and by competitive protein binding assays (Eisman
15 et al., Anal. Biochem. 80: 298-305 (1977); and Haddad et al., J. Clin. Endocr. 33: 992-995 (1971)). For example, the vitamin D transport protein known as DBP, which has a strong preference for binding 25-OH-D was used in the competitive binding assay (Bouillion et al., J. Steroid
20 Biochem. 13: 1029-1034 (1980)).

Also, various competitive binding assays have been employed to assay for the presence of $1,25-(OH)_2-D$. (Shigeharu et al., Anal. Biochem. 116: 211-222 (1981); Eisman et al., Arch. Biochem. Biophys. 176: 235-243
25 (1976); Perry et al., Biochem. Biophys. Res. Comm. 112: 431-436 (1983); Bouillion et al., Ann. Endocrin. 41: 435-436 (1980); Bouillion, Clin. Chem. 26: 562-567 (1980); Bouillion, Eur. J. Biochem. 66: 285-291 (1976)). In such assays, vitamin D and its metabolites were extracted from
30 blood serum or plasma with an organic solvent. The extract was then purified by column chromatography and HPLC to yield $1,25-(OH)_2-D$. The purified $1,25-(OH)_2-D$

was then measured.

In general, most assays presently being employed in the art to determine the concentration of vitamin D are heterogeneous assays. Furthermore, most of these assays
5 rely on the addition of an organic solvent to release the vitamin D and/or vitamin D metabolites from the binding proteins, for example DBP. (Schmidt-Gayk, Dynamics of Bone and Cartilage Metabolism, Chapter 26, Table V). That is, the addition of organic solvent to the samples
10 causes denaturation and precipitation of the serum proteins including DBP, and subsequently, the precipitated protein can be spun out of solution and the released metabolites remain in solution in the organic layer. This organic layer containing the released
15 metabolite is then extracted and transferred into another system for analysis.

Although the present existing assay systems are useful, their reliance on an organic solvent to release the vitamin D and/or vitamin D metabolite is problematic.
20 The problems with using an organic solvent include: (A) difficulties in the handling of volatile organic solvents, (B) laborious manual extractions, (C) loss of patient identification since two transfer steps are required, and (D) loss of precision in measurement.

25 The loss in precision may be caused by the heterogeneous extraction step, since heterogeneous extraction can be very technique dependent. For example, in a heterogeneous extraction step, some matrix components may be extracted along with the organic layer
30 containing the metabolite. One such matrix component is lipid, which has been shown to cause an elevated measurement of the metabolite values.

There is a need to have improved kits and methods for determining the concentration of vitamin D and its metabolites in a body fluid of a mammal.

5 Summary of the Invention

In accordance with the present invention, a kit is featured for determining the concentration of a vitamin D component. In a broad embodiment, the kit comprises a releasing composition. The releasing composition
10 facilitates in releasing the vitamin D component from a vitamin D component binding-protein. In one embodiment, the releasing composition is substantially free of an organic solvent. In one embodiment, the kit further comprises a detecting composition. The detecting
15 composition facilitates in determining the concentration of the vitamin D component.

Further in accordance with the present invention, a kit according to this invention may be useful for determining the concentration of the vitamin D component
20 present in a mammal fluid. The mammal fluid may be milk, whole blood, serum, plasma and mixtures thereof. For example, a mammal fluid may comprise a human serum.

Still further in accordance with the present invention, the vitamin D component may be a vitamin D, a
25 precursor of a vitamin D, a metabolite of a vitamin D and mixtures thereof. For example, a vitamin D component comprises a 25-OH-D.

Still further in accordance with the present invention, the releasing composition comprises an aqueous
30 base component, for example, NaOH or KOH. In one embodiment, the releasing composition comprises about 0.1 to about 1.0 M of the aqueous base component. In one

embodiment, the releasing composition is substantially free of an organic solvent. In one embodiment, the releasing composition may be adapted to various bio-assay systems to assay for a vitamin D component.

5 Still further in accordance with the present invention, the releasing composition further comprises a cyclo-oligomer component, for example a cyclodextrin, derivatives thereof and mixtures thereof. In one embodiment, the releasing composition comprises about
10 0.01 to about 5% of the cyclo-oligomer component.

Still further in accordance with the present invention, the releasing component further comprises about 0.5 to about 5% of a metal salicylate, including sodium salicylate.

15 Still further in accordance with the present invention, the releasing component further comprises about 0.01 to about 0.1% of a surfactant.

Still further in accordance with the present invention, the releasing composition forms a homogeneous
20 mixture with a mammal fluid.

Still further in accordance with the present invention, the detecting composition comprises a host component and a partner component, wherein the host component binds to the partner component to form a
25 partner/host complex. The concentration of the complex should be proportional to the concentration of the vitamin D component. In one embodiment, the concentration of the complex is inversely proportional to the concentration of the vitamin D component. In one
30 embodiment, the host component comprises an antibody and/or portions thereof.

Still further in accordance with the present

invention, the host component is labeled. In one embodiment, the host component is labeled with a chemiluminescent label, for example acridinium, a fluorescent label and/or a radio-active label.

5 Still further in accordance with the present invention, the partner component comprises a vitamin D component linked to a separator component, wherein the separator component is a solid phase, such as antibody or a magnetic particle.

10 Still further in accordance with the present invention, the partner component binds to the host component through at least one intermediate binding component, for example, a vitamin D binding-protein.

Still further in accordance with the present
15 invention, a method is featured for assaying a body fluid sample for the concentration of a vitamin D component. The method comprises the steps of releasing the vitamin D component from the vitamin D component binding-protein by contacting the sample with a releasing
20 composition in a holder, and determining the concentration of the vitamin D component.

Still further in accordance with the present invention, the determining step includes the steps of (a) adding a detecting composition to the holder, the
25 detecting composition comprises a host component and a partner component, the host component binds to the partner component to form a partner/host complex, (b) isolate the complex in the tube, (c) measuring the concentration of the complex by measuring the
30 concentration of the host component in the complex, the concentration of the complex is proportional to the concentration of the vitamin D component.

Any feature or combination of features described herein are included within the scope of the present invention provided that the features included in any such combination are not mutually inconsistent as will be
5 apparent from the context, this specification, and the knowledge of one of ordinary skill in the art.

Additional advantages and aspects of the present invention are apparent in the following detailed description and claims.

10

Detailed Description of the Invention

In a broad embodiment, the present invention provides a kit for determining the concentration of a vitamin D component. In one embodiment, the kit, is
15 useful for determining the concentration of a vitamin D component found in a mammal, for example a human. In a preferred embodiment, the kit is useful for determining the concentration of a vitamin D component in a body fluid of a mammal. The body fluid may include, without
20 limitation, milk, whole blood, serum, plasma or mixtures thereof. Preferably, the body fluid is serum of a human.

A vitamin D component includes a vitamin D, vitamin D precursors and vitamin metabolites. Without intending to be limited by the examples, a vitamin D component may
25 include, for example, vitamin D₂, D₃, D₄, D₅, and D₆; precursors of a vitamin D such as 7-dehydrocholesterol; and metabolites of vitamin D such as metabolites of vitamin D₂, D₃, D₄, D₅, and D₆, preferably 1,25-OH-D₂, and more preferably 25-OH-D. Often, a vitamin D component is
30 bound to a vitamin D component binding protein. For example, in human serum, 25-OH-D is bound to a protein; this protein is commonly referred to in the art as a DBP.

A kit of the present invention comprises a releasing composition. The releasing composition of this invention is adaptable for use in or with various bio-assay techniques known in the art to determine the concentration of a vitamin D component. Preferably, the releasing composition of the present invention substantially eliminates the need to purify the vitamin D component prior to determining its concentration. Also preferably, a bio-assay kit comprising a releasing composition of this invention has improved precision in the measurement of a vitamin D component.

In one embodiment, the releasing composition comprises an aqueous base component and facilitates in releasing the vitamin D component from a binding entity. The binding entity includes, without limitation, a binding protein or a binding lipid. The aqueous base component comprises any molecule which is effective in raising and maintaining the pH of the releasing composition to a pH of greater than 7, preferably greater than 8, more preferably about 12 to about 14, even more preferably about 13. In one embodiment, the base component comprises a metal hydroxide, for example a NaOH and/or a KOH. In one embodiment, the releasing composition is substantially free of an organic solvent, for example acetonitrile.

In a preferred embodiment, the releasing composition comprises about 0.1 to about 1 M of the aqueous base component. More preferably, the releasing composition comprises about 0.35 to about 0.5 M of the aqueous base component. For example, a releasing composition in accordance with the present invention may include about 0.35 to about 0.5 M of NaOH.

Without wishing to limit the invention to any particular theory or mechanism of operation, it is believed that the aqueous base component denatures, preferably irreversibly denatures, the vitamin D binding protein, i.e. DBP, and causes the bound vitamin D component to be released. Additionally, the aqueous base component may also irreversibly denature other plasma or serum proteins, such as albumin. It is further believed that these proteins may interfere in the detection of the vitamin D component and their denaturation reduces this effect.

In one embodiment, the releasing composition further comprises a cyclo-oligomer component. In a preferred embodiment, the cyclo-oligomer component is at least effective in sequestering interfering components. Interfering components, without limitation, include hydrophobic molecules, such as lipids. Interfering components interfere with the measurement of the vitamin D component, and therefore, its sequestration substantially reduces interference of the measurement. For example, lipids have been shown to erroneously increase the measurement values of a vitamin D component.

In a preferred embodiment, the cyclo-oligomer component comprises an α -cyclodextrin. In a more preferred embodiment, the cyclo-oligomer component comprises a β -randomly methylated cyclodextrin. The cyclo-oligomer component may also include, without limitation, derivatives of α -cyclodextrin, β -cyclodextrin, derivatives of β -cyclodextrin, γ -cyclodextrin, derivatives of γ -cyclodextrin, carboxymethyl- β -cyclodextrin, carboxymethyl-ethyl- β -

cyclodextrin, diethyl- β -cyclodextrin, dimethyl- β -cyclodextrin, methyl- β -cyclodextrin, random methyl- β -cyclodextrin, glucosyl- β -cyclodextrin, maltosyl- β -cyclodextrin, hydroxyethyl- β -cyclodextrin, hydroxypropyl-
5 β -cyclodextrin, sulfobutylether- β -cyclodextrin, and the like and mixtures thereof. As used herein, the term "derivatives" as it relates to a cyclodextrin means any substituted or otherwise modified compound, which has the characteristic chemical structure of the parent
10 cyclodextrin.

In one embodiment, the releasing composition comprises about 0.01 to about 5% of the cyclo-oligomer component. For example, in a preferred embodiment, the releasing composition comprises about 2% of an α -
15 cyclodextrin. In another preferred embodiment, the releasing composition comprises about 0.05% of a β -randomly methylated cyclodextrin. In an alternative embodiment, the releasing composition is substantially free of a cyclo-oligomer component.

20 In one embodiment, the releasing component further comprises a salicylate and the like, such as 8-anilino-1-naphthalenesulfonic acid (ANS). Without wishing to be limited by any theory or mechanism of operation, it is believed that the salicylate or the like of the present
25 invention is at least effective to release the vitamin D component from binding to proteins and/or solubilize the vitamin D component in the releasing composition. In a preferred embodiment, the salicylate is a metal salicylate, for example sodium salicylate. The release
30 composition preferably comprises about 0.5 to about 5% of a metal salicylate, for example a sodium salicylate. In

an alternative embodiment, the releasing composition is substantially free of a salicylate and the like.

In one embodiment, the releasing component further comprises a surfactant and the like. Without wishing to
5 be limited by any theory or mechanism of operation, it is believed that the surfactant and the like of the present invention is at least effective to block the vitamin D component from being attached to lipids, proteins and the like. In a preferred embodiment, the surfactant is
10 tween-20 or triton X-100. The releasing composition of the present invention may include about 0.01 to about 0.1% of a surfactant. For example, 0.05% of tween-20 may preferably be employed in the present invention. In another preferred embodiment, the releasing component is
15 substantially free of a surfactant.

The releasing composition of the present invention may comprise an aqueous base component and may be substantially free of the cyclo-oligomer component, the salicylate and the surfactant. In a preferred
20 embodiment, the release component comprises an aqueous base component and at least a cyclo-oligomer component, a salicylate and/or a surfactant. In a more preferred embodiment, the release component comprises an aqueous base component and a cyclo-oligomer component. In an
25 even more preferred embodiment, the release component comprises an aqueous base component, a cyclo-oligomer component and a salicylate. For example, a preferred releasing composition comprises about 0.1 to about 1 M of an aqueous base component, about 0.01 to about 5% of a
30 cyclo-oligomer component and about 0.01 to about 5% of a salicylate.

As discussed above, the releasing composition of this invention is adaptable for use in or with various bio-assay techniques known in the art to determine the concentration of a vitamin D component. For example, in
5 one embodiment, a kit of the present invention comprises a releasing composition and a detecting composition.

In a broad embodiment, the detecting composition of the present invention comprises a host component. Preferably, the detecting composition further comprises a
10 partner component. The host component is any molecule or set of molecules capable of selectively binding to a partner component or a vitamin D component. For example, a host component may be an aptamer or a molecular imprint polymer. In one embodiment, a host component is an
15 antibody, or a portion of an antibody, for example a Fab portion, capable of binding to a partner component. The antibody of the present invention may be monoclonal or polyclonal.

Various methods are known in the art to produce an
20 antibody specific toward a certain antigen, for example a partner component. For example, an antibody may be raised from a rabbit injected with an antigen, the antigen being the partner component or a part thereof. Additionally, synthetic antibodies may also be made.
25 (See U.S. Patent No. 5,110,833, the disclosure of which is incorporated in its entirety by reference herein.)

The specific composition of the partner component depends on the nature of the assay. For example, if a competitive assay technique is employed, the partner
30 component may comprise a competing molecule which is capable of competing with the released vitamin D component at the binding site of the host component to

form a partner/host complex. Preferably this competing molecule is linked to a separator component, wherein the separator component eventually facilitates in separating out and isolating the partner/host complex. The
5 separator component may, without limitation, comprise a peptide, an antibody, a small molecule, a polymer particle and/or a magnetic particle. The competing molecule may be linked to the separating component by various means known in the art. For example, the
10 molecule may be linked to the separator component through a chemical bond or via an antibody.

In one embodiment, the separator component may be isolated from solution, along with anything else which is presently attached to it, for example a competing
15 molecule, by various means. For example, if the separating component is an antibody, an Enzyme-Linked-ImmunoSorbent Assay (ELISA) technique or the like may be employed to isolate the separating component. If the separating component is a magnetic particle, then a
20 magnetic field may be employed to isolate the separating component from solution.

In one embodiment, the partner component comprises a vitamin D component linked to a solid phase, for example a particle, preferably a magnetic particle. In a
25 preferred embodiment, the vitamin D component directly links to a particle. For example, a 3-hydroxyl of the 25-OH-D (a vitamin D component) may form a direct covalent bond with a functional group present on a particle; such group includes, without limitation, -COOH,
30 -NH₂, epoxide, tosyl, and -SH. In a more preferred embodiment, the vitamin D component links to a particle through at least one linking component, for example a

biotin, derivatives thereof and the like. (see Holick et al U.S. Patent No. 5,981,779, the disclosure of which is incorporated in its entirety herein by reference). For example, in one embodiment, the partner component
5 comprises a 25-OH-D linked to a magnetic particle via a biotin.

In one embodiment, a partner component competes with the vitamin D component, for example, the released 25-OH-D, to bind to the host component, for example an
10 antibody, to form a partner/host complex. The partner/host is then isolated and its concentration is determined. The concentration of the vitamin D component, for example the 25-OH-D, should be directly proportional to the concentration of the partner/host
15 complex.

In an alternative embodiment, the partner component comprising a competing molecule linked to a separator component competes with the vitamin D component to bind to one or more intermediate binding components. At least
20 one of the intermediate binding components has a binding site for a vitamin D component. In this case, the host component can form a complex with the partner component through an intermediate binding component. (When the partner component forms an a complex with a host
25 component through an intermediate component, a partner/intermediate/host complex is formed. The partner/intermediate/host complex may sometimes be referred to herein as the "partner/host" complex for clarity.) For example, in one embodiment, the partner
30 component comprises a 25-OH-D linked to a magnetic particle through a biotin, and this partner component competes with a released 25-OH-D at the binding site of

an intermediate binding component, for example a DBP. After the partner component binds to the intermediate binding component, the host component, for example an antibody, may form a complex with the partner component
5 through the intermediate binding component, by binding to the intermediate component. In a preferred embodiment, the host component antibody which binds to a DBP (hereinafter "anti-DBP") is a rabbit antibody purchased under the trade name Rabbit Anti-Human Gc-Globulin from
10 DAKO in Denmark.

Furthermore, the host component may be present when the partner component competes with the vitamin D component to bind to the intermediate binding component; or the partner/intermediate complex may be first isolated
15 from solution and the host component is subsequently introduced to form a partner/intermediate/host complex.

The concentration of the partner/host complex formed in accordance with this invention should be proportional to the concentration of the vitamin D component being
20 measured. (Herein after and throughout "partner/host complex" may refer to a partner/host complex or partner/intermediate/host complex.) In one embodiment, the concentration of the partner/host complex may be measured by isolating the partner/host complex and
25 measuring the level of the host component and/or measuring the level of the intermediate component if the partner component forms a complex with the host component through an intermediate component. Various methods known in the art may be adapted and employed to measure the
30 concentration of the host component and/or the intermediate component.

For example, in one embodiment, the host component

and/or the intermediate component (hereinafter "signal compounds") may be labeled. Preferably, the signal compound is labeled at a position where the signal compound may still bind to a partner component and the label may still be detected. Labels which may be employed in accordance with this invention include, without limitation, chemiluminescent molecules, fluorescent molecules, enzymes, co-enzymes, isotopes and sensitizers.

Chemiluminescent molecules useful in this invention include, without limitation, luciferin, luminol, pyrogallol, isoluminol, aequorin, cyclic arylhydrazides, dioxetanes, rhodium chelates (electrochemiluminescent), oxalate esters, thermochemiluminescent labels, acridinium and the like. These labels may be attached to a protein, for example an anti-DBP, with techniques well known in the art. (See U.S. Patent No. 5,284,952, the disclosure of which is incorporated in its entirety herein by reference.) In one embodiment, a signal compound such as an anti-DBP may be labeled with an acridinium by employing the methods found in U.S. Patent No. 5,284,952. (See also U.S. Patent Nos. 5,110,932 and 5,338,847, the disclosures of which are incorporated in their entirety herein by reference.) In this embodiment, the acridinium labels the lysine residue of the anti-DBP. It has been shown that a sufficient number of the anti-DBP is labeled at the appropriate places to be effective as a labeled host component.

Fluorescent molecules useful in this invention include, without limitation, umbelliferone, fluorescein, rhodamine, Texas red dyes, phthalocyanines, coumarin, squaraine, anthracene, erythrosine, europium chelates and

the like. For example, a signal compound such as a DBP may be labeled with an umbelliferone. (See also, U.S. Pat. No. 3,901,654, the disclosure of which is incorporated in its entirety by reference herein.)

5 Enzymes useful in this invention include enzymes capable of catalyzing a reactant to produce a product, wherein the product can be detected. Exemplary enzymes which have been developed and used in an assay are those described in U.S. Pat. Nos. 3,654,090; 3,791,932;
10 3,839,153; 3,850,752; U.S. Pat. No. 3,817,837; 3,879,262; Journal of Immunological Methods 1: 247(1972); and the Journal of Immunology 109:129(1972), the disclosures of which are incorporated in their entirety herein by reference. Other enzymes which may be used in accordance
15 with this invention includes galactosidase, gluconidase, phosphatase, peptidase, alkaline phosphatase and the like. In one embodiment, an enzyme in accordance with this invention may be able to catalyze a chemiluminescent reactant, wherein one of the resulting product is a light
20 signal, which can be detected.

Co-enzymes useful in this invention include molecules and/or proteins which facilitate an enzyme to catalyze a reactant to produce a detectable product, for example light. A co-enzyme may include, without
25 limitation, FAD and NAD. In one embodiment, an anti-DBP may be labeled with a NAD. (See for example, U.S. Patent 4,380,580, the disclosure of which is incorporated in its entirety herein by reference.)

Isotopes useful in this invention include, without
30 limitation, H^3 , P^{32} and F^{18} . Other labels may include a non-active precursor of a spectrophotometrically-active substance (British Pat. No. 1,392,403 and French Pat. No.

2,201,299, which patents correspond to U.S. Pat. No. 3,880,934) and electron spin resonance moieties (U.S. Pat. No. 3,850,578).

Sensitizers useful in this invention include, without limitation, those disclosed in U.S. Patent No. 5,705,622, the disclosure of which is incorporated in its entirety herein by reference. For example, an acceptor molecule may be attached to a DBP and an acceptor molecule may be attached to a vitamin D component.

Various kits comprising a releasing composition and a detecting composition as disclosed in this invention may be used for determining the concentration of a vitamin D component. However, such are only examples of the many possible kits which may employ a releasing composition to assay for a vitamin D component. Other kits are contemplated to be within the scope of this invention. For example, the following patents, articles and instruction manuals disclose assay methods which may be adapted to include a releasing composition of the present invention: U.S. Patent No. 4,935,339 (discloses a delayed capture assay); No. 4,121,1978; No. 5,232,836; No. 5,064,770; No. 5,202,266; No. 4,816,417; No. 5,821,020 and No. 5,981,779; Competitive Protein-Binding Radioassays for 25-OH-D by Haddad; Instruction Manual of Jummun Diagnostik Entitled 25-OH Vitamin D Enzyme-based Protein-Binding-Assay; Instruction Manual of Incstar Catalog No. 68100; Directional insert of Nichols Institute entitled 25-OH-D 60T Kit; Directional Insert of BioSource Europe S.A. entitled 25OH-VIT.D3-RIA-CT; Directional Insert of IDS, Inc. entitled Gamma-B25-Hydroxy Vitamin D RIA AA-35PS (Jan 6, 1999, issue 3); Journal of Pharmaceutical and Biomedical Analysis

(1999); Determination of Vitamin D3 metabolites/state-of-the-art by Luque de Castro, et al.; Clin. Chem. 30/11, 1731-1736 (1984); Two Direct (Nonchromatographic) Assays for 25-hydroxyvitamin D by Bouillon, et al. The disclosures of which are incorporated in their entirety by reference herein. One adaptation would be to replace the compositions used therein with the present releasing composition.

Without limitation, Table I identifies some general assays (A through H) in which a releasing composition of this invention may be employed. Assay A has been discussed above.

Assay B may similarly employ a releasing composition to release the vitamin D component of a body fluid sample. Once released, it may be added to a composition or a series of compositions comprising a labeled DBP and a vitamin D component linked to a separator component, for example a magnetic particle. Preferably, the released vitamin D component does not have to be purified prior to adding to said composition. In other words, a composition comprising a labeled DBP and a vitamin D component linked to a separator component may be added to the homogeneous mixture of body fluid sample and releasing composition.

Assays C through G may similarly employ a releasing composition to release the vitamin D component from a body fluid sample. Once released, a detecting composition or series of detecting compositions, comprising the components identified in the respective assays, may be added. In assay G, the conditions conducive for the binding of the biotin to the avidin may be similar to that disclosed in U.S. Patent No.

5,395,938, (Title: Biotinylated Chemiluminescent Labels and Their Conjugates, Assays and Assay Kits), the disclosure of which is incorporated in its entirety herein by reference.

5 Assay H may also employ a releasing composition of the present invention to release the vitamin D component from a body fluid sample. After it is released the vitamin D component is allowed to bind directly to a host component, for example an antibody. The released vitamin
10. D component is also allowed to bind to a labeled antibody. Preferably the host component and the labeled antibody are able to bind to different determinants of the released vitamin D component. (See U.S. Patent No. 5,641,690, which discloses immunometric assay methods,
15 the disclosures of which are incorporated in its entirety herein by reference). The host/vitamin-D/labeled antibody complex may be isolated and counted utilizing techniques commonly known in the art, for example ELISA.

Table I




Table I ContinuedDefinitions

A vitamin D component added to the assay system to compete with the released vitamin D component.



A vitamin D component released by the releasing composition.



A vitamin D binding protein, for example DBP.



A solid phase, for example, a magnetic particle.



A host component, for example an antibody, which recognizes a vitamin D component.



A molecule which can bind to a biotin, for example streptavidin or avidin.



A label, for example acridinium.



A biotin.



A protein, for example an antibody, capable of binding to a vitamin D component at a site other than the one that is recognized by a host component.



Means that one entity is linked to another entity. The linkage may be direct, or through another molecule or molecules.



Means that one entity is linked to another entity. The linkage may be direct, or through another molecule or molecules.

In a broad embodiment, the kit of the present invention may be adapted to be employed in an automated assay system to determine the concentration of a vitamin
5 D component. For example, the kit of the present invention may preferably be used in conjunction with the Nichols Advantage system.

EXAMPLE 1

10 An automated method for detecting a 25-OH-D

Currently, immunoassays measuring the concentration of 25-OH-D use organic solvents to release the 25-OH-D. The addition of an organic solvent to the serum results in either a precipitate or a two-layer liquid.
15 Thereafter, the layer containing the released 25-OH-D is (manually) separated and is mixed with another composition to measure the concentration of the 25-OH-D. The step of separating the 25-OH-D is (a) time consuming, and (b) a source of error in the assays, especially if
20 the separation step is performed manually.

The present invention eliminates the need to purify or substantially purify the released 25-OH-D prior to determining its concentration and allows for automation of the assay. In this example, the automated release of
25 the 25-OH-D from a binding protein and determination of the 25-OH-D concentration are performed on the Nichols Advantage system. This process is wholly automated. Manual pipetting and centrifugation are not needed. The assay kit use in conjunction with the Nichols Advantage
30 System includes a release composition and detecting composition.

Releasing Composition

The releasing composition comprises an aqueous base component (0.1-1.0 M NaOH), beta-randomly methylated cyclodextrin (0.01-5%), and sodium salicylate (0.01-5%).
5 The pH of the releasing composition is about 13.

Detecting Composition

The detecting composition comprises a magnetic particle suspension with 25-OH-D immobilized on the
10 surface, a DBP, an anti-DBP and stabilizers. Also, the detecting composition has a pH of about 6 to about 9, preferably 8.3. Furthermore, the detecting composition also has a buffer capacity to bring the pH of the total mixture (the homogeneous mixture and the detecting
15 composition) into the range where the DBP is stable and capable of binding 25-OH-D. Preferably the total mixture has a pH of about 6 to about 9, preferably 8.3. The buffer system used comprises 200 mM TRIS, 150 mM NaCl, 0.9% gelatin, 1% bovine serum albumin and 0.09% sodium
20 azide.

The streptavidin coated particles are about 0.5-5 microns in size, and may be purchased from Seradyne, Dynal, Bangs Laboratories. The 25-OH-D is attached to the particle via a biotin or the like (Holick et al U.S.
25 Patent No. 5,981,779, the disclosure of which is incorporated in its entirety herein by reference). The amount of particles used in the assay is about 5 to about 30 uL (preferably 25 uL) of a 1.25-2.5 mg/mL suspension.

The concentration of the DBP is in the range about
30 0.2 to about 1.0 ug/mL and is added to the assay in the amount of about 10 to about 280 uL, preferably about 250 uL.

The anti-DBP, preferably purchased from DAKO, is labeled with an acridinium sulfonyl chloride derivative, a chemiluminescent label. Preferably, the acridinium derivative is labeled by a method disclosed in U.S. Patent No. 5,284,952, the disclosure of which is incorporated in its entirety herein by reference. In brief, the conjugate is prepared by adding the reactive label in acetonitrile to the antibody in carbonate buffer at pH 9.6. The reaction proceeds at ambient temperature for a period of about half an hour and the unattached label is removed by gel filtration chromatography. The labeled anti-DBP is diluted to a concentration of about 0.2-10 ug/mL in buffer with a pH of about 6.0. The amount of acridinium labeled anti-DBP that is added to the assay is about 10 to about 280 uL, preferably about 50uL.

In preferred embodiments, a stabilizer may be added to a composition of the assay. For example, a stabilizer may be added to the detecting composition. Stabilizers include gelatin, bovine serum albumin, egg albumin, polyethylene glycol, poly vinyl alcohol, poly vinyl pyrrolidone and the like and mixtures thereof. These stabilizers may be effective in blocking of the non-specific binding of the assay components.

About 10-100uL, preferably 20uL, of a patient body fluid sample and about 10-100uL, preferably 60 uL, of the releasing composition are pipetted into a holder, for example a cuvette well, forming a homogeneous mixture. The releasing composition acts on the patient sample in an incubator at a constant temperature (about 25-37°C, preferably 37 °C) for a specified time (1-60 min., preferably 21 min.). When the release is complete, a

detecting component is pipetted directly into the homogeneous mixture, forming a total mixture. The total mixture is incubated for about 30 min to about 60 min., preferably about 42 min. During this incubation period
5 the 25-OH-D-biotin-magnetic-particle competes with the 25-OH-D of the patient's sample to form a complex with the labeled-DBP. Subsequently, the 25-OH-D-biotin-magnetic-particle/labeled-DBP complex is isolated by a wash. The concentration of the complex is determined by
10 a chemiluminescent reading. The concentration of the complex is inversely proportional to the concentration of the 25-OH-D concentration in the patient's sample.

In addition to decreasing the labor requirements, this assay method allows for positive patient
15 identification. Positive patient identification can be achieved because the Nichols Advantage system pipettes the detecting composition directly into the sample holder containing the releasing composition and the body fluid, and every sample holder has a bar code to which the
20 result matches.

EXAMPLE 2

Precision of assays: Intra CV

Intra Coefficient of Variation (CV) studies for the
25 determination of the 25-OH-D were conducted with various systems, including a system using a kit of the present invention in conjunction with Nichols automated assay machine, the Nichols Advantage. The studies using the kit of the present invention and Nichols Advantage
30 automated technique involves obtaining a patient sample and performing the assay 20 times. The results are shown in Table II.

Table II

	Dose (ng/ml)	CVD
Nichols Advantage Automated assay Run #1	12.3	7.7%
	34.6	3.5%
	86.0	2.0%
Nichols Advantage Automated assay Run #2	9.5	14.1%
	46.8	2.6%
	58.6	2.6%
¹²⁵ I Assay Kit #1	Dose	CVD
	26.5	5.3%
	58.4	5.0%
	151.0	6.1%
¹²⁵ I Assay Kit #2	Dose	CVD
	8.6	11.7%
	22.7	10.5%
	33.0	8.6%
	49.0	12.5%

Each patient sample was assayed 20 times. Of the 20 results, a mean and a standard of deviation were determined. $CV = (\text{standard of deviation}/\text{mean}) \times 100$. CVD is CV of Dose, wherein Dose is the concentration of the 25-OH-D. The lower the CV, the better the precision of measurement. For example, a good CVD is <8%.

For each Dose range, the CVDs of the assays using an assay kit of the present invention in conjunction with the Nichols Advantage system are better than other assay systems. For example, the assay kit of the present invention has a CVD of 2.6% at a Dose of 46.8; the ^{125}I Assay Kit #2 has a CVD of 12.5% at a Dose of 49.0; and the ^{125}I Assay Kit #1 has a CVD of 5.0% at a Dose of 58.4.

15

EXAMPLE 3

Luminol Chemiluminescent Direct Labeling of DBP

About 1 mg of Vitamin D binding protein, DBP, is buffer exchanged three times with 100 mM PBS at pH 8.2. The final volume was 200 uL. Succinyl-amino-buty-ethyl-isoluminol NHS ester (4.6 mg) is dissolved in DMF. The isoluminol (17 uL) is added to the DBP and the mixture is maintained at room temperature for about 1 hour with occasional shaking. The unreacted label is removed by size exclusion chromatography using Sephadex-G25 (23 X 1 cm) and 100 mM PBS, pH 6.0 as the mobile phase.

25

EXAMPLE 4

Direct Chemiluminescent Direct Labeling of DBP

About 1 mg of Vitamin D binding protein, DBP, is buffer exchanged three times with 20 mM sodium carbonate buffer at pH 9.6. The final volume is 500 uL. Two

30

aliquots of fluorescein-5-isothiocyanate (FITC) are dissolved in DMF just prior to addition to give a concentration of 1.25 mg/mL. The first FITC aliquot (5 uL) is added and the mixture is allowed to react at room temperature. After 15 minutes the second aliquot (5 uL) is added and the reaction proceeds for an additional 15 minutes. The unreacted label is removed by size exclusion chromatography using Sephadex G-25 (2 cm X 33 cm) and 100 mM, pH 7.4 mobile phase.

10

EXAMPLE 5

Biotin Direct Labeling of DBP for "Upside-Down" Assay

About 2.0 mg of DBP is buffer exchanged with 20 mM bicarbonate at pH 9.6 three times to give a final volume of 1.0 mL. Two aliquots of sulfo-NHS-LC-biotin (Pierce) are dissolved in water to give a 1 mg/mL solution just prior to addition. The first aliquot (10 uL) is added and the reaction is maintained at room temperature for 5 minutes. The second aliquot (10 uL) is added and the reaction proceeds for an additional 4 minutes. The unreacted biotin is removed by size exclusion chromatography using Sephadex G-25 and 100 mM PBS, pH 7.4 as the mobile phase.

25

EXAMPLE 6

Acridinium Direct Labeling of anti-25-Hydroxyvitamin D monoclonal antibody

2.5 mg of antibody is buffer exchanged into 20 mM bicarbonate buffer, pH 9.6, three times to give a final volume of 1.75 mL. Sulfonylchloride acridinium ester is dissolved in sufficient acetonitrile to give a 1.75 mg/mL solution. The acridinium ester (52.5 uL) is added to the

antibody and the reaction mixture is maintained at room temperature about 0.5 hour. The unreacted label is removed by size exclusion chromatography using Sephadex, G-75 and 100 mM PBS, pH 6.0 as the mobile phase.

5

EXAMPLE 7

Acridinium Indirect Labeling

Anti-DBP (Dako) is buffer exchanged to yield 2.2 mg of protein in 1.1 mL of pH 9.6, 20 mM bicarbonate buffer.

10 A solution of sulfonylchloride acridinium ester (1.6 mg/mL) is in acetonitrile. The acridinium (4.4 uL) is added to the antibody and the reaction mixture is maintained at room temperature for 15 minutes. A second aliquot of acridinium (4.4) uL is added for and

15 additional 15 minutes. The unreacted label is removed by chromatography on a mixed bed column (Sephadex 6B/Sephadex G-75, 1.5 cm X 48 cm).

EXAMPLE 8

Direct labeling of DBP with HRP

20 Activated peroxidase from Boehringer Mannheim Biochimica is used for labeling. The DBP (4 mg) is dissolved in pH 9.8, 100 mM carbonate buffer (1 mL). This solution is added to the POD (8 mg). The reaction

25 proceeded at room temperature for 1 hour. The conjugation is quenched by addition of 120 uL of triethanolamine solution (pH 8, 30%). Sodium borohydride (150 uL) freshly dissolved in water (8 mg/mL) is added to the solution maintained at room temperature for about

30 0.5 hour. Additional sodium borohydride (75 uL) is provided and the reaction proceed for an additional 2 hours at room temperature. The conjugate is dialyzed

with pH 6.0 PBS with glycine.

While this invention has been described with respect to various specific examples and embodiments, it is to be understood that the invention is not limited thereto and
5 that it can be variously practiced with the scope of the following claims.

What is claimed is:

1. A kit for determining a concentration of a vitamin D component comprising:

5

a releasing composition; and

a detecting composition,

10 the releasing composition comprises an aqueous base component and facilitates in releasing the vitamin D component from a vitamin D component binding-protein, the detecting composition facilitates in determining the concentration of the vitamin D component.

15

2. A kit of claim 1 being useful for determining the concentration of the vitamin D component present in a mammal fluid.

20 3. A kit of claim 1 wherein the mammal fluid is selected from the group consisting of milk, whole blood, serum, plasma and mixtures thereof.

25 4. A kit of claim 1 wherein the mammal fluid comprises a human serum.

5. A kit of claim 1 wherein the vitamin D component is selected from the group consisting of a metabolite of vitamin D₂, D₃, D₄, D₅, and D₆.

30

6. A kit of claim 1 wherein the vitamin D component comprises a 25-OH-D.

7. A kit of claim 1 wherein the vitamin D component comprises a 1, 25-(OH)₂-D.
- 5 8. A kit of claim 1 wherein the aqueous base component comprises NaOH.
9. A kit of claim 1 wherein the aqueous base component comprises KOH.
- 10 10. A kit of claim 1 wherein the releasing composition comprises about 0.1 to about 1.0 M of the aqueous base component.
- 15 11. A kit of claim 1 wherein the releasing composition comprises about 0.35 to about 0.5 M of the aqueous base component, wherein the aqueous base component is NaOH.
12. A kit of claim 1 wherein the releasing composition
20 is substantially free of an organic solvent.
13. A kit of claim 1 wherein the releasing composition further comprises a cyclo-oligomer component.
- 25 14. A kit of claim 13 wherein the cyclo-oligomer component comprises a cyclodextrin.
15. A kit of claim 13 wherein the cyclo-oligomer component is selected from the group consisting of alpha-
30 cyclodextrin and beta-randomly methylated cyclodextrin.
16. A kit of claim 13 wherein the releasing composition

comprises about 0.01 to about 5% of the cyclo-oligomer component.

17. A kit of claim 13 wherein the releasing component
5 comprises about 2% of the cyclo-oligomer component,
wherein the cyclo-oligomer component is an alpha-cyclodextrin.

18. A kit of claim 13 wherein the releasing component
10 comprises about 0.05% of the cyclo-oligomer component,
wherein the cyclo-oligomer component is a beta-randomly methylated cyclodextrin.

19. A kit of claim 1 wherein the releasing component
15 further comprises about 0.5 to about 5% of a metal salicylate, including sodium salicylate.

20. A kit of claim 1 wherein the releasing component
further comprises about 0.01 to about 0.1% of a
20 surfactant.

21. A kit of claim 20 wherein the surfactant is selected from the group consisting of tween-20 and triton X-100.

22. A kit of claim 1 wherein the releasing composition
25 forms a homogeneous mixture with a mammal fluid.

23. A kit of claim 1 wherein the releasing composition
comprises

30

about 0.1 to about 1.0 M of an aqueous base component;

about 0.01 to about 5% of a cyclo-oligomer component; and

5 about 0.01 to about 5% of a metal salicylate.

24. A kit of claim 23 wherein the aqueous base component is NaOH, the cyclo-oligomer component is cyclodextrin and the metal salicylate is sodium salicylate.

10

25. A kit of claim 1 wherein the detecting composition comprises a host component and a partner component, wherein the host component binds to the partner component to form a partner/host complex, the concentration of the
15 complex is proportional to the concentration of the vitamin D component.

26. A kit of claim 25 wherein the host component comprises an antibody, portions thereof, or mixtures
20 thereof.

27. A kit of claim 25 wherein the host component is labeled with a chemiluminescent label, a fluorescent label or a radio-active label.

25

28. A kit of claim 25 wherein the host component is an antibody labeled with acridinium.

29. A kit of claim 25 wherein the partner component
30 comprises a vitamin D component linked to a separator component, the separator component is a solid phase or an antibody.

30. A kit of claim 29 wherein the separator component comprises a magnetic particle.
- 5 31. A kit of claim 29 wherein the partner component comprises a vitamin D component linked to a magnetic particle.
- 10 32. A kit of claim 31 wherein the partner component competes with the vitamin D component to bind to the host component.
- 15 33. A kit of claim 32 wherein the host component is an antibody labeled with acridinium.
34. A kit of claim 32 wherein the partner component binds to the host component through at least one intermediate binding component.
- 20 35. A kit of claim 34 wherein the intermediate component is labeled.
36. A kit of claim 34 wherein the intermediate component is labeled and the host component is not labeled.
- 25 37. A kit of claim 34 wherein at least one intermediate binding component comprise a vitamin D binding-protein.
- 30 38. A kit of claim 25 wherein the partner component competes with a vitamin D component to form a complex with the host component, the partner component comprises a vitamin D component linked to a magnetic particle, the

partner component binds to the host component through a vitamin D binding-protein, the host component comprises an antibody labeled with acridinium.

5 39. A kit of claim 38 wherein the concentration of the complex is inversely proportional to the concentration of the vitamin D component.

40. A kit of claim 1 wherein the concentration of the
10 vitamin D component is determined with a higher precision than that of an assay kit relying on an organic solvent, to release the vitamin D component from the binding-protein.

15 41. A kit of claim 1, wherein the releasing composition forms a homogeneous mixture with a body fluid containing the vitamin D component.

42. A kit for determining a concentration of a vitamin D
20 component comprising:

a releasing composition comprising about 0.1 to about 1.0 M of NaOH or KOH, 0 to about 5% of a cyclodextrin, 0 to about 5% of salicylate and 0 to about
25 0.1% of a surfactant,

a detecting composition comprising an antibody labeled with acridinium and a partner component, wherein the partner component competes with a vitamin D component
30 to form a complex with the antibody, the partner component comprises a vitamin D component linked to a magnetic particle, the partner component binds to the

antibody through a vitamin D binding-protein.

43. A kit of claim 42 wherein the releasing composition comprises about 0.35 to about 0.5 M of NaOH, about 2% of
5 alpha-cyclodextrin and about 2% of salicylate, the releasing composition being substantially free of an organic solvent.

44. A releasing composition comprising an aqueous base
10 component, wherein the releasing composition facilitates in releasing a vitamin D component from a binding-protein, the releasing composition being useful in determining the concentration of a vitamin D component.

45. A releasing composition of claim 44 wherein the
15 aqueous base component comprises NaOH.

46. A releasing composition of claim 44 wherein the
aqueous base component comprises KOH.

20

47. A releasing composition of claim 44 comprising about
0.1 to about 1.0 M of the aqueous base component.

48. A releasing composition of claim 44 comprising about
25 0.35 to about 0.5 M of the aqueous base component, wherein the aqueous base component is NaOH.

49. A releasing composition of claim 44 being
substantially free of an organic solvent.

30

50. A releasing composition of claim 44 further
comprising a cyclo-oligomer component.

51. A releasing composition of claim 50 wherein the cyclo-oligomer component comprises a cyclodextrin.
- 5 52. A releasing composition of claim 50 wherein the cyclo-oligomer component is selected from the group consisting of alpha-cyclodextrin and beta-randomly methylated cyclodextrin.
- 10 53. A releasing composition of claim 50 comprising about 0.01 to about 5% of the cyclo-oligomer component.
54. A releasing composition of claim 50 comprising about 2% of the cyclo-oligomer component, wherein the cyclo-
15 oligomer component is an alpha-cyclodextrin.
55. A releasing composition of claim 50 comprising about 0.05% of the cyclo-oligomer component, wherein the cyclo-oligomer component is a beta-randomly methylated
20 cyclodextrin.
56. A releasing composition of claim 44 further comprising about 0.5 to about 5% of a metal salicylate, including sodium salicylate.
- 25 57. A releasing composition of claim 44 further comprising about 0.01 to about 0.1% of a surfactant.
58. A releasing composition of claim 57 wherein the
30 surfactant is selected from the group consisting of tween-20 and triton X-100.

59. A releasing composition of claim 44 wherein the releasing composition forms a homogeneous mixture with a body fluid sample containing the vitamin D component.

5 60. A releasing composition of claim 59 wherein a detecting composition is added to the homogeneous mixture to determine the concentration of the vitamin D component.

10 61. A releasing composition of claim 44 being employed in a biochemical assay.

62. A releasing composition of claim 61 wherein the biochemical assay is a homogeneous biochemical assay.

15

63. A releasing composition comprising

about 0.1 to about 1.0 M of an aqueous base component;

20

about 0.01 to about 5% of a cyclo-oligomer component; and

25 about 0.01 to about 5% of a metal salicylate, the releasing composition is useful in determining the concentration of a vitamin D component.

64. A releasing component of claim 63 wherein the aqueous base component is NaOH, the cyclo-oligomer component is cyclodextrin and the metal salicylate is sodium salicylate.

30

65. A releasing component of claim 63 being substantially free of an organic solvent.

66. A method of assaying a body fluid sample for the concentration of a vitamin D component, the method comprising the steps of:

releasing the vitamin D component from the vitamin D component binding-protein by contacting the sample with a releasing composition in a holder; and

determining the concentration of the vitamin D component.

67. A method of claim 66 wherein the vitamin D is released into a homogeneous mixture of the body fluid sample and the releasing composition.

68. A method of claim 66 wherein the releasing composition comprises

about 0.1 to about 1.0 M of an aqueous base component;

0 to about 5% of a cyclo-oligomer component;

0 to about 5% of a metal salicylate; and

0 to about 0.1% of a surfactant.

69. A method of claim 68 wherein the aqueous base component is NaOH, the cyclo-oligomer component is

cyclodextrin, the metal salicylate is sodium salicylate and the surfactant is tween-20.

70. A method of claim 66 wherein the releasing
5 composition is substantially free of an organic solvent,
including an organic solvent.

71. A method of claim 67 wherein the determining step
includes the steps of:

10

adding a detecting composition to the holder, the
detecting composition comprises a host component and a
partner component, the host component binds to the
partner component to form a partner/host complex;

15

isolate the complex in the tube;

measuring the concentration of the complex by
measuring the concentration of the host component in the
20 complex, the concentration of the complex is proportional
to the concentration of the vitamin D component.

72. A method of claim 71 wherein the host component is
an antibody labeled with acridinium.

25

73. A method of claim 72 wherein the concentration of
the host is measured by detecting the emitted level of
chemiluminescence.

30 74. A method of claim 71 wherein the partner component
competes with a vitamin D component to form a complex
with the host component, the partner component comprises

a vitamin D component linked to a magnetic particle, the partner component binds to the host component through a vitamin D binding-protein.

- 5 75. A method of claim 66 wherein the determining step includes the steps of:

adding a partner component and a vitamin D binding-protein to the tube, the partner component competes with
10 the vitamin D component to bind to the vitamin D component binding-protein to form a partner/binding-protein complex;

isolate the partner/binding-protein complex in the
15 tube; and

adding a host component, the host component binds to the partner/binding-protein complex to form a partner/binding-protein/host component complex.

20

measuring the concentration of the partner/binding-protein/host component complex by measuring the concentration of the host component in the complex, the concentration of the complex is proportional to the
25 concentration of the vitamin D component.

76. A method of claim 75 wherein the host component is an antibody labeled with acridinium.

- 30 77. A method of claim 75 wherein the concentration of the host is measured by detecting the emitted level of chemiluminescence.

78. A method of claim 75 wherein the partner component comprises a vitamin D component linked to a magnetic particle.

5

79. A method of claim 66 providing a more precise determination of the vitamin D component as compared to method using a releasing composition comprising an organic solvent.

10

80. A method of assaying a body fluid sample for the concentration of a 25-OH-D component, the method comprising the steps of:

15 releasing the 25-OH-D from the 25-OH-D binding-protein by contacting the sample with a releasing composition in a holder, including a cuvette, the releasing composition comprises about 0.1 to about 1.0 M of an aqueous base component, 0 to about 5% of a cyclo-
20 oligomer component, 0 to about 5% of a metal salicylate, and 0 to about 0.1% of a surfactant; and

adding a detecting composition to the holder,

25 the detecting composition comprises an antibody labeled with acridinium, a 25-OH-D binding-protein and a partner component, the partner component comprises a 25-OH-D linked to a magnetic particle,

30 the partner component competes with the released 25-OH-D to bind to the 25-OH-D binding-protein to form a partner/binding-protein complex,

the antibody binds to the partner/binding-protein complex to form an antibody/binding-protein/partner component complex;

5

isolating the antibody/binding-protein/partner component complex in the tube; and

measuring the concentration of the antibody/binding-protein/partner component complex by measuring the concentration of the labeled antibody in the complex, the concentration of the complex is proportional to the concentration of the vitamin D component.

10 81. A method of claim 80 wherein the concentration of the labeled antibody is measured by detecting the emitted level of chemiluminescence.

82. A method of claim 80 wherein the releasing composition comprises about 0.35 to about 0.5 M of NaOH, about 2% of alpha-cyclodextrin and about 2% of salicylate.

83. A method of claim 80 wherein the releasing composition is substantially free of an organic solvent, including an organic solvent.

84. A method of assaying a body fluid sample for the concentration of a vitamin D component, the method comprising the steps of:

forming a homogeneous mixture of a body fluid sample

containing the vitamin D component and a releasing component; and

adding a detecting composition to the mixture to
5 determine the concentration of the vitamin D component.